

## ***Cryptococcus neoformans*, Isolate 10**

**Catalog No. NR-41300**

**For research use only. Not for human use.**

### **Contributor:**

Qiang Qiang Zhang, Professor, Huashan Hospital, Fudan University, Shanghai, China

### **Manufacturer:**

BEI Resources

### **Product Description:**

Classification: *Filobasidiaceae*, *Cryptococcus*

Species: *Cryptococcus neoformans*

Isolate: 10

Original Source: *Cryptococcus neoformans* (*C. neoformans*), isolate 10 was obtained from human cerebrospinal fluid in China in May 2011.<sup>1</sup>

The *Cryptococcus* species complex is currently composed of two species, *C. neoformans* and *C. gattii*. These species are best recognized as the agents of cryptococcosis, an AIDS-defining illness.<sup>2,3</sup> *C. neoformans* is divided into two varieties, *C. neoformans* var. *grubii* (serotype A) and *C. neoformans* var. *neoformans* (serotype D).<sup>2</sup> In the current classification scheme, there are five distinct lineages, referred to as VNI, VNII, VNB, VNIII and VNIV.<sup>2,3</sup> The two varieties (*neoformans* and *grubii*) are able to recombine and produce diploid or aneuploid intervarietal AD hybrids.<sup>2</sup> *C. neoformans* has been widely associated with avian excreta.<sup>2</sup>

### **Material Provided:**

Each vial of NR-41300 contains approximately 0.4 mL of yeast culture in 20% glycerol.

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

### **Packaging/Storage:**

NR-41300 was packaged aseptically in cryovials and is provided frozen on dry ice. The product should be stored at -60°C or colder. For long term storage, cryogenic temperature (-130°C or colder), preferably in the vapor phase of a liquid nitrogen freezer, is recommended.

### **Growth Conditions:**

#### Media:

Yeast Mold broth or equivalent

Yeast Mold agar or equivalent

#### Incubation:

Temperature: 25°C to 30°C

Atmosphere: Aerobic

### Propagation:

1. Keep vial frozen until ready for use; thaw rapidly in a waterbath at 25°C to 30°C. Typically, this takes less than 5 minutes.
2. Immediately after thawing, inoculate an agar plate with approximately 40 µL of thawed culture or transfer the entire thawed aliquot into a single tube of broth.
3. Incubate the plate or tube at 25°C to 30°C for 2 to 4 days.

### **Citation:**

Acknowledgment for publications should read "The following reagent was obtained through BEI Resources, NIAID, NIH: *Cryptococcus neoformans*, Isolate 10, NR-41300."

### **Biosafety Level: 2**

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. Biosafety in Microbiological and Biomedical Laboratories. 5th ed. Washington, DC: U.S. Government Printing Office, 2009; see [www.cdc.gov/biosafety/publications/bmbl5/index.htm](http://www.cdc.gov/biosafety/publications/bmbl5/index.htm).

### **Disclaimers:**

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at [www.beiresources.org](http://www.beiresources.org).

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

### **Use Restrictions:**

**This material is distributed for internal research, non-commercial purposes only.** This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals

contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a license is required. U.S. Government contractors may need a license before first commercial sale.

**References:**

1. Zhang, Q. Q., Personal Communication.
2. Cogliati, M. "Global Molecular Epidemiology of *Cryptococcus neoformans* and *Cryptococcus gattii*: An Atlas of the Molecular Types." *Scientifica (Cairo)* 2013 (2013): 675213. PubMed: 24278784.
3. Zhu, P., et al. "Congenic Strains for Genetic Analysis of Virulence Traits in *Cryptococcus gattii*." *Infect. Immun.* 81 (2013): 2616-2625. PubMed: 23670558.

ATCC<sup>®</sup> is a trademark of the American Type Culture Collection.

