

***Bacillus anthracis*, Strain Sterne ΔGBAA0419-2**

Catalog No. NR-10001

Product Description: *Bacillus anthracis* (*B. anthracis*), strain Sterne ΔGBAA0419-2 is a deletion mutant of the toxigenic acapsulate original Sterne strain (34F2), constructed by replacing codons 10 through 14 with three in-frame stop codons followed by the recognition site for *Bam*HI (to facilitate screening of the correct mutation). The remainder of the putative calcium/proton exchanger gene (GBAA0419-2) retains the wild type sequence.

Lot¹: 58441522

Manufacturing Date: 17DEC2008

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis Cellular morphology Colony morphology ² Sporulation Motility β-hemolysis Capsule (India ink staining) Tenacious Analytical profile index (API [®] 50 CHB including API [®] 20E; ONPG to GEL only) Nitrate reduction FAME analysis	Gram-positive rod Report results Positive Non-motile Non-hemolytic Negative Positive Consistent with <i>B. anthracis</i> Positive Consistent with <i>B. anthracis</i>	Gram-positive rod Circular, flat, erose, ground-glass, opaque and grey (Figure 1) Positive Non-motile Non-hemolytic Negative Positive Consistent with <i>B. anthracis</i> Positive Consistent with <i>B. anthracis</i>
Genotypic Analysis Sequencing of 16S ribosomal RNA gene (1435 base pairs)	Consistent with <i>Bacillus cereus</i> group	Consistent with <i>Bacillus cereus</i> group ^{3,4}
PCR Assay of Extracted DNA 16S ribosomal RNA gene Presence of virulence plasmids pXO1 (<i>aat</i>) pXO2 (<i>at</i> , <i>capA</i> , <i>capB</i> , <i>capC</i>)	~ 1500 bp amplicon ~ 125 bp amplicon No amplicons	~ 1500 bp amplicon ~ 125 bp amplicon No amplicons
Viability (post-vialing)⁵	Growth	Growth

¹*B. anthracis*, strain Sterne ΔGBAA0419-2 was deposited by Philip C. Hanna, Associate Professor, Department of Microbiology and Immunology, University of Michigan Medical School, Ann Arbor, Michigan. NR-10001 was produced by inoculation of the deposited material into Tryptic Soy Broth and grown 24 hours at 37°C. Broth inoculum was added to Kolles which were grown 24 hours at 37°C to produce this lot.


²24 hours at 37°C on Tryptic Soy Agar with 5% sheep blood

³*Bacillus cereus* group species (*B. cereus*, *B. thuringiensis*, *B. mycoides*, and *B. anthracis*) cannot be classified based on 16S sequence (Spencer, R. C. "Bacillus anthracis." *J. Clin. Pathol.* 56 (2003): 182-187. PubMed: 12610093).

⁴Also consistent with *Bacillus subtilis*

⁵24 hours at 37°C in Tryptic Soy Broth

Figure 1



Date: 06 AUG 2009

Signature: Signature on File

Title: Technical Manager, BEI Authentication or designee

ATCC[®], on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC[®]'s knowledge.

ATCC[®] is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

