

Cryptococcus gattii, Strain R265

Catalog No. NR-50184

Product Description: *Cryptococcus gattii* (*C. gattii*), strain R265 was isolated from a human on Vancouver Island, Canada during the outbreak that began in the late 1990's.

Lot¹: 63779296

Manufacturing Date: 09OCT2015

TEST	SPECIFICATIONS	RESULTS
Phenotypic Analysis		
Cellular morphology ²	Report results	Globose, single or budding (Figure 1A)
Colony morphology ²	Report results	Butyrous, circular, smooth, entire and off-white (Figure 1B)
Canavanine-glycine-bromthymol blue (CGB) differential medium ³	Blue (<i>C. gattii</i>)	Blue (<i>C. gattii</i>)
Genotypic Analysis		
Sequencing of partial 18S rRNA gene, internal transcribed spacer (ITS) 1, 5.8S rRNA gene, ITS 2, partial 28S rRNA (~ 520 base pairs)	≥99% sequence identity to <i>C. gattii</i> (GenBank: FJ914888.1)	100% sequence identity to <i>C. gattii</i> (GenBank: FJ914888.1)
Sequencing of 26S rRNA gene (620 base pairs)	≥99% sequence identity to <i>C. gattii</i> (GenBank: KC171326.1)	100% sequence identity to <i>C. gattii</i> (GenBank: KC171326.1)
Purity⁴		
Nutrient broth with 0.1% Yeast Extract at 25°C	No bacterial growth	No bacterial growth
Nutrient broth with 0.1% Yeast Extract at 37°C	No bacterial growth	No bacterial growth
Viability (post-freeze)²	Growth	Growth


¹NR-50184, lot 63779296, was produced by incubation of seed material on modified Sabouraud Dextrose medium and incubated for 3 days at 25°C in an aerobic atmosphere. Yeast were harvested from agar plates with 20% glycerol prior to vialing.

²3 days at 25°C in an aerobic atmosphere on modified Sabouraud Dextrose agar

³3 days at 27°C in an aerobic atmosphere. CGB medium differentiates *C. gattii* from *C. neoformans* based on the ability of *C. gattii* isolates to grow in the presence of L-canavanine and to assimilate glycine as a sole carbon source, resulting in a blue color. *C. neoformans* isolates will show yellow to light-green on CGB medium. [McTaggart, L., et al. "Rapid Identification of *Cryptococcus neoformans* var. *grubii*, *C. neoformans* var. *neoformans*, and *C. gattii* by Use of Rapid Biochemical Tests, Differential Media, and DNA Sequencing." *J. Clin. Microbiol.* 2011 (49): 2522-2527. PubMed: 21593254.]

⁴Clarity of broth was determined by visual inspection after 3 days at 25°C and 37°C in an aerobic atmosphere.

Figure 1: Cellular and Colony Morphology



Certificate of Analysis for NR-50184

Date: 15 JUL 2016

Signature:



BEI Resources Authentication

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

