

***Coccidioides immitis*, Strain 2102**

**Catalog No. NR-48936**

**Product Description:** *Coccidioides immitis* (*C. immitis*), strain 2102 was isolated in the 1990s from a human in San Diego, California, USA.

**Lot<sup>1</sup>: 70003192**

**Manufacturing Date: 17APR2017**

TEST	SPECIFICATIONS	RESULTS
<b>Phenotypic Analysis</b> Colony morphology <sup>2</sup>	Report results	Circular, raised, undulate, membranous, rough and white (Figure 1)
<b>Genotypic Analysis<sup>3</sup></b> Sequencing of partial 18S ribosomal RNA (rRNA) gene, internal transcribed spacer (ITS) 1, 5.8S rRNA gene, ITS 2, partial 26S rRNA (~ 500 base pairs) Sequencing of 28S rRNA gene (~ 620 base pairs)	≥ 99% sequence identity to <i>C. immitis</i> , strain 2394 (GenBank: AATX01000460.1) ≥ 99% sequence identity to <i>C. immitis</i> , strain 2394 (GenBank: AATX01000460.1)	99.2% sequence identity to <i>C. immitis</i> , strain 2394 (GenBank: AATX01000460.1) <sup>4</sup> 99.8% sequence identity to <i>C. immitis</i> , strain 2394 (GenBank: AATX01000460.1) <sup>4</sup>
<b>Purity<sup>5</sup></b> Nutrient broth with 0.1% Yeast Extract at 25°C Nutrient broth with 0.1% Yeast Extract at 37°C	No bacterial growth No bacterial growth	Non-turbid growth Non-turbid growth
<b>Viability (post-freeze)<sup>2</sup></b>	Growth	Growth

<sup>1</sup>NR-48936 was produced by inoculation of the deposited material into Emmon's Modified Sabouraud Dextrose broth and grown for 13 days at 37°C in an aerobic atmosphere with 5% CO<sub>2</sub>. The broth growth was added to Emmon's Modified Sabouraud Dextrose kolles and grown for 12 days at 37°C at 37°C in an aerobic atmosphere with 5% CO<sub>2</sub>. The agar growth was harvested from the kolles with 20% glycerol to produce this lot.


<sup>2</sup>6 days at 37°C in an aerobic atmosphere with 5% CO<sub>2</sub> on Emmon's Modified Sabouraud Dextrose agar

<sup>3</sup>*C. immitis* and *C. posadasii* cannot be identified to the species level based on ITS and large subunit rRNA sequences (Tintelnot, K., et al. "Taxonomic and Diagnostic Markers for Identification of *Coccidioides immitis* and *Coccidioides posadasii*." *Med. Mycol.* 45 (2007): 385-393. PubMed: 17654264.).

<sup>4</sup>Also consistent with *C. posadasii*

<sup>5</sup>Clarity of broth was determined by visual inspection after 6 days in an aerobic atmosphere with 5% CO<sub>2</sub>.

**Figure 1: Colony Morphology**



/Heather Couch/

Heather Couch

23 AUG 2018

Program Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

